

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Genetic variability detected at the lactoferrin locus (LTF) in the Italian Mediterranean river buffalo**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1530132> since 2018-03-15T23:11:01Z

*Published version:*

DOI:10.1071/AN14612

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*[Animal Production Science, published on line on 18<sup>th</sup> February 2015, DOI*

*http://dx.doi.org/10.1071/AN14612]*

*ovvero [Pauciullo A. et al., CSIRO Publishing, 2015, pagg. 1-6]*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*[<http://www.publish.csiro.au/paper/AN14612.htm>]*

**Genetic variability detected at the lactoferrin locus (*LTF*) in the Italian Mediterranean river buffalo**

<sup>A</sup>Alfredo Pauciullo, <sup>B, E</sup>Luigi Ramunno, <sup>C</sup>Nicolò P. P. Macciotta, <sup>C</sup>Giustino Gaspa, <sup>D</sup>Angelo Coletta, <sup>B</sup>Elisa Apicella, <sup>B</sup>Daniela Gallo, <sup>B</sup>Gianfranco Cosenza

Short title: Genetic variability at the river buffalo *LTF* gene

<sup>A</sup> *ISPAAM, Laboratory of Animal Cytogenetics and Gene Mapping, National Research Council, via Argine 1085, 80147 Napoli, Italy*

<sup>B</sup> *Department of Agriculture, University of Naples Federico II, via Università 100, 80055 Portici, Napoli, Italy*

<sup>C</sup> *Department of Agriculture, University of Sassari, via De Nicola 9, Sassari, 07100 Italy*

<sup>D</sup> *Italian National Association of Buffalo Breeders (ANASB), Via F. Petrarca 42-44 Località Centurano, Caserta, 81100 Italy*

<sup>E</sup> *Corrisponding author: L. Ramunno*

*Department of Agriculture, University of Naples Federico II, via Università 100, 80055 Portici, Napoli, Italy*

e-mail: [ramunno@unina.it](mailto:ramunno@unina.it)

**Abstract.** Lactoferrin (*LTF*) is multi-functional protein belonging to the whey protein fractions of the milk. The gene *LTF* encoding for such protein is considered a potential candidate for body measurement, milk composition and yield. This study reports on the genetic variability at *LTF* locus in the Italian Mediterranean river buffalo and its possible association with milk yield. Eleven polymorphic sites were found in the DNA fragment spanning the exons 15-16. In particular, the intron 15 was extremely polymorphic with 9 SNPs detected, whereas the remaining 2 SNPs were exonic mutations (g.88G>A at the exon 15 and g.1351G>A at the exon 16) and both synonymous. The genotyping of the informative samples evidenced 3 haplotypes, whose frequencies were 0.6; 0.3 and 0.1 respectively, whereas the analysis of the exonic SNPs showed a perfect condition of linkage disequilibrium (g.88A/g.1351G and g.88G/g.1351A). The association study carried out by using the SNP g.88G>A showed that buffalo *LTF* gene has no statistically significant influence on daily milk yield. This study adds knowledge to the genetic variability of a species less investigated than the other ruminant species, that may serve as a useful tool for large-scale screening of buffalo populations.

**Additional keywords:** *Bubalus bubalis*, lactoferrin, polymorphism, haplotype, milk yield, marker assisted selection

## 1 **Introduction**

2 In Italy there is a buffalo industry of limited size, in comparison with many east Asian countries,  
3 but of great economic relevance due to the production of Mozzarella PDO (Protected  
4 Denomination of Origin - Reg. EC 510/2006) one of the most popular and spread Italian dairy  
5 product. River buffaloes farmed in Italy are named ‘Mediterranean Italian’, aiming at  
6 distinguishing them from other European populations which are not at the same genetic level  
7 (Borghese, 2005).

8 In 2012, the consistency of the Italian buffalo stock was about 387,000 heads, with an  
9 increase of 7.2% over the previous year. The production of buffalo milk in 2012 amounted to  
10 1,924,553 tons, with an increase of 7.79% compared to 2010 (<http://www.aia.it>). Also an  
11 improvement in buffalo milk composition occurred in the last few years, with the average protein  
12 and fat content moving from 4.65% and 8.10% in 2003 to 4.70% and 8.30% in 2012,  
13 respectively, (<http://www.anasb.it/home.htm>). These figures could be ascribed to the effort that  
14 has been undertaken in improving the buffalo management system, feeding and also the breeding  
15 programme.

16 Advances of molecular genetics offer the possibility to investigate genomic regions that  
17 affect traits of economic importance and to identify useful genetic polymorphisms for marker-  
18 assisted selection (MAS) programmes. In recent years, some studies have been carried out on  
19 buffalo for the identification of genetic polymorphisms at milk protein *loci* (Masina *et al.* 2007;  
20 Cosenza *et al.* 2009a, b; Pauciullo *et al.* 2011; Cosenza *et al.* 2014), and candidate genes  
21 responsible for the variation of the quali-quantitative characteristics of the Mediterranean water  
22 buffalo milk were identified (Cosenza *et al.* 2007; Pauciullo *et al.* 2010a), as well as significant  
23 associations with milk yield (Pauciullo *et al.* 2012a, b) and milk coagulation properties (Bonfatti  
24 *et al.* 2012) have been found.

Lactoferrin (*LTF*) is an iron-binding glycoprotein and it belongs to the whey protein fractions of the milk. It is a multi-functional protein playing an important role in antibacterial, antiviral, antifungal, anti-inflammatory, anti-oxidant, immune-modulatory activities and in the regulation of iron metabolism (Vogel *et al.* 2002; Kang *et al.* 2008). Apart from all the aforementioned functions, the efficient expression of lactoferrin as milk component is also very important. In fact, *LTF* is considered a potential candidate gene for body measurement (withers height, body length and chest circumference) as well as for milk composition and yield (Guo *et al.* 2010). Lactoferrin concentration is quite variable across different species and in particular in buffalo's milk ( $0.332 \pm 0.165$  g/kg) appeared to be lower than that reported in human milk, but higher than found in bovine milk (Giacinti *et al.* 2013).

The *LTF* gene has been particularly investigated in cattle and goat and many polymorphisms have been found in these two species (Seyfert *et al.* 1994; Lee *et al.* 1997; Li *et al.* 2004; Daly *et al.* 2006; Kaminski *et al.* 2006; Kang *et al.* 2008; Arnould *et al.* 2009; O'Halloran *et al.* 2009; Huang *et al.* 2010; Pauciullo *et al.* 2010b), some of which showed associations with milk production traits and antimicrobial properties (Lee *et al.* 1997; Li *et al.* 2004; Guo *et al.* 2010). In contrast, few studies have been carried out in buffalo. Currently, only short, partial genomic or cDNA sequences, all related to Indian buffaloes, are available in EMBL. Two synonymous SNPs (adenine-guanine transitions) within the exon 16 of *LTF* gene and different polymorphisms in non-coding regions (intron 4 and intron 9) have been found by comparing lactoferrin sequences of different Indian buffalo breeds (Murrah, Jaffarabadi, Toda, Marathwada, Pandharpuri and Mehsana) (Kathiravan *et al.* 2009; Kathiravan *et al.* 2010). Polymorphisms within the promoter region have been also found in different Khuzestan buffalo breeds (Khatibi *et al.* 2013). In contrast, so far no sequence and no polymorphisms are available for the Italian Mediterranean buffalo.

Considering the split architecture of the *LTF* locus (17 exons over ~30kb in cattle), the aim of our work was to sequence part of this gene in the Italian Mediterranean river buffalo, to detect genetic variability at the *LTF* gene and to investigate possible associations with milk yield.

## **Materials and methods**

### *Sample collection*

Individual blood samples were collected from 553 Italian river buffaloes belonging to 11 farms, located in Salerno and Caserta province (Southern Italy). Sampling was carried out in collaboration with the Italian National Association of Buffalo Breeders (ANASB).

### *DNA isolation and PCR amplification conditions*

DNA was isolated from leukocytes, using the procedure described by Goossens and Kan (1981). DNA concentration and OD<sub>260/280</sub> ratio of the samples were measured by the Nanodrop ND-2000C Spectrophotometer (Thermo Scientific).

The DNA regions of the *LTF* gene spanning from the 15<sup>th</sup> exon to the 16<sup>th</sup> exon, including the flanking regions, of ten individual samples randomly chosen were amplified by iCycler (BioRad) using the following primers: (forward) LTF15F: 5'-GCTGATGCAGCCTTCTCT-3'; (reverse) LTF16R: 5'-TTTAAACCCACATCACCCCT-3'. The sequences of the two primers correspond to nucleotides 15432-15449 and to the complementary nt 17054-17073, respectively, of the partial sequence of the bovine gene deposited at EMBL (acc. no [Z93399](#)).

The 25 µl reaction mix comprised 100 ng of genomic DNA, 50 mM KCl, 10 mM Tris-HCl (pH 9.0), 0.1% Triton X-100, 3 mM MgCl<sub>2</sub>, 200 nmol of each primer, dNTPs each at 400 mM, 2.5 U Taq DNA Polymerase (Promega, Madison, WI), and 0.04% BSA. The amplification program consisted of an initial denaturation at 97°C for 2 min, annealing of primers at 62°C for

45 s, and an extension step at 72°C for 2 min; then 30 cycles of denaturation at 94°C for 45 s, annealing at 62°C for 45 s, and extension at 72°C for 2 min (except for the final extension of 10 min). PCR products were analysed directly by electrophoresis in 1.5% TBE agarose gel (Bio-Rad, CA, USA) in 0.5X TBE buffer and stained with SYBR<sup>®</sup> green nucleic acid stain (Lonza Rockland, Inc, USA). PCR products were sequenced on both strands at CEINGE - Biotecnologie Avanzate (Naples, Italy).

#### *Bioinformatic and statistical analysis*

Allelic frequencies and Hardy-Weinberg equilibrium ( $\chi^2$  test) were calculated for each SNP. Homology searches, comparison among sequences, and multiple alignments were accomplished using DNAsis-Pro (Hitachi Software Engineering Co., Japan).

The intron 15 was analyzed for potential microRNA sequence by using the bovine miRBase database (<http://www.mirbase.org/search.shtml>), whereas the program TFSEARCH (<http://www.rwcp.or.jp/papia/> Heinemeyer *et al.* 1998) was used to identify potential transcription factor binding sites and TargetScanHuman ([http://www.targetscan.org/vert\\_61/](http://www.targetscan.org/vert_61/)) was used to identify microRNA target regions.

Measures of linkage disequilibrium ( $D'$  and  $r^2$ ) were estimated using Haploview software *ver.* 4.2 (<http://www.broadinstitute.org/haploview/haploview>). Haplotype structure was defined according to Gabriel *et al.* (2002).

A total of 7,260 records for milk yield measured monthly by the official recording system of the Italian Association of Buffalo Breeders (ANASB) on 1,097 lactations of the 553 buffaloes were used. Data were collected in the period January 2009-September 2012.

Associations between *LTF* polymorphism and milk yield was investigated with the following mixed linear model (SAS Institute, Cary NC, USA):



$$y_{ijklm} = \mu + lc_i + hys_j + DIM_k + LTF_l + DIM_k * LTF_l + a_m(LTF_l) + e_{ijklm} \quad (1)$$

where

$y_{ijklm}$  = record of milk yield;

$\mu$  = overall mean;

$lc_i$  = fixed effect of the  $i^{th}$  age-parity, ( $i=1, \dots, 56$ );

$hys_j$  = fixed effect of the  $j^{th}$  heard-year-season, ( $j=1, \dots, 24$ );

$DIM_k$  = fixed effect of the  $k^{th}$  month of lactation, ( $k=1, \dots, 10$ );

$LTF_l$  = fixed effect of the  $l^{th}$  LTF genotype, ( $l=AA, AG, GG$ );

$a_m(LTF_l)$  = random effects of individual cow ( $m=1, \dots, 553$ ) nested within LTF genotype;

$e_{ijklm}$  = random residual.

Pairwise comparisons among different levels of fixed effects included in model were performed using a Bonferroni adjusted test.

## Results and discussion

### *Polymorphism detection*

In order to detect polymorphisms at the Mediterranean river buffalo *LTF* locus, the DNA region spanning from the last 2 nucleotides of the 14<sup>th</sup> intron to the first 2 nucleotides of the 16<sup>th</sup> intron was sequenced for a total of 1505 bp (EMBL acc. no. [HG515533](#)).

The DNA fragment sequenced showed a homology of 95, 93 and 91% respectively with cattle (EMBL acc. no. [Z933399](#)), goat (EMBL acc. no. [FJ609300](#)) and sheep (EMBL acc. no. [FJ541507](#)) *LTF* gene. All splice junctions follow the 5'GT/3'AG splice rule.

1       The analysis and the comparison of the obtained sequences showed a total of 11  
2 polymorphic sites. Nine of these were localized at the level of the 15<sup>th</sup> intron (g.299C>T,  
3 g.367A>G, g.893A>G, g.1051A>G, g.1110A>G, g.1113A>G, g.1120C>T, g.1127C>T,  
4 g.1214C>T), none of which affected canonical splicing sites.

5       Moreover, we detected one SNP in exon 15 (g.88G>A) and another in exon 16  
6 (g.1351G>A). Both the exonic mutations were found to be synonymous with no changes in the  
7 amino acid sequence: p.Thr582 and p.Lys628 respectively. The enumeration of amino acids starts  
8 at the first amino acid of the mature peptide, with number 1 (EMBL acc. no. [AJ005203](#)).

9       The presence of the guanine in position 88 characterizes also other *LTF* sequences of  
10 buffalo (EMBL acc. no. [JF825526](#), [AJ005203](#)) and of other ruminants, such as goat (EMBL acc.  
11 no. [X78902](#), [DQ387456](#), [CHU53857](#)), sheep (EMBL acc. no. [NM\\_001024862](#), [KC161426](#)) and  
12 cattle (EMBL acc. no. [GQ351344](#), [FJ589071](#)), therefore its presence might be indicative of an  
13 ancestral condition. Analogously, the presence of guanine in position 1,351 characterizes the  
14 sequences relating to other ruminant and non-ruminant species, except for some breeds of India  
15 (EMBL acc. no. [JF825526](#), [AJ005203](#), [EU518482](#)) that is characterized by the presence of  
16 adenine. However, the same locus was described as polymorphic by Kathiravan *et al.* (2010)  
17 which also report a G>A transition. Therefore, this finding seems to confirm the guanine as  
18 possible ancestral condition.

19       A comparison of the sequence of the 15<sup>th</sup> and 16<sup>th</sup> exons (185 bp and 191 bp, respectively)  
20 of the gene that we sequenced with the published sequence of the buffalo *LTF* cDNA (EMBL  
21 acc. no. [AJ005203](#)) shows two nucleotide differences located at 15<sup>th</sup> exon, none of which results  
22 in an amino acid substitution: g.58C>T, p.Leu572 and g.140C>T, p.Leu600

23       The sequence analysis for the informative samples showed a condition of linkage  
24 disequilibrium for most of the detected polymorphic sites. Only three haplotypes were found: H1

(ATGGAAGCCCG), H2 (GCAAGGACCCA) and H3 (GCGGGGATTTA) whose frequencies were 0.6, 0.3 and 0.1 respectively (Fig. 1). (Insert Fig.1 here)

#### *Intron 15 analysis for potential microRNA sequence*

The analysis of the intron 15 (1,125 bp) showed two mature sequence for the miR-181a (EMBL: HG515533, nucleotides 889-906) and miR-2459 (nucleotides 1076-1094). MicroRNA (miR) are small noncoding RNA that regulate gene expression post-transcriptionally and play a key role in development and specific biological processes. The expression pattern of the miR-181a was already investigated in the bovine mammary gland, however no significant difference were reported at different stages of the lactation cycle (Wang *et al.* 2012). On contrary, miR-2459 expression pattern in mammary gland has not been investigated yet. Both microRNA are involved in the innate and adaptive immune response regulating target genes like TRAF6 (TNF receptor associated factor 6), TRAF3 (TNF receptor associated factor 3), TNF, MMD (monocyte to macrophage differentiation-associated), etc... and their presence in *LTF* gene might be connected with the defense mechanism related to this milk protein.

The analysis of the regulatory regions upstream and downstream of both miR sequences did not evidence any influence of the detected polymorphisms on putative consensus sequences for transcription factors. However the SNP g.893A>G fell within the mature sequence of the miR-181. Nevertheless, except when conserved sequences in splice sites are changed, the real effects of non-coding polymorphisms cannot be easily predicted.

#### *Genotyping of Mediterranean river buffalo LTF alleles*

Although the detected polymorphisms in the coding region do not result in amino-acid changes, they can still affect gene function by altering the stability, splicing or localization of the mRNA.

In general, synonymous changes are less likely to be associated with functional changes or diseases, and so should be given lower priority for genotyping. Nevertheless, because of their potential effect on mRNA stability, they should have higher priority than polymorphisms that lie deep within introns (Risch 2000). For these reasons we decided to genotype the investigated buffalo population only for exonic SNPs: g.88G>A and g.1351G>A.

The genotyping of 553 buffalo DNA samples was performed at the KBiosciences (Herts, UK, <http://www.kbioscience.co.uk>) laboratory. The major allele had a relative frequency of about 0.682 in both *loci* and  $\chi^2$  values showed that there was no evidence of departure from the Hardy-Weinberg equilibrium ( $P \leq 0.05$ ). A complete linkage disequilibrium was observed for these 2 SNPs (average  $D'$  and  $r^2$  values were 1.0 and 1.0 respectively), whose haplotype (g.88A/g.1351G and g.88G/g.1351A) frequencies were 0.318 and 0.682, respectively, i.e. equal to allelic ones. According to Gabriel *et al.* (2002), *loci* can be considered as being in the same haplotype block. Thus only one SNP (g.88G>A) was considered in running model (1). Genotype distribution of buffalo cows for the SNP g.88G>A are reported in Table 1. (Insert table 1 here)

#### *Association of LTF polymorphism on milk yield*

Despite the relatively large amount of data analyzed and the accurate measures on milk yield, the present study suggests that no direct effect of the investigated *LTF* polymorphisms on the production trait studied. The larger difference in the least squares means for milk yield was estimated in 0.36 kg/d between the genotypes AA and AG, whereas the GG showed an average milk production of 8.35 kg/d. These differences were not significant, therefore this SNP appears to be selectively neutral in relation to milk yield. Although not significant, this result adds new information to the recent reports on polymorphisms found in other genes influencing the milk yield in river buffalo (Table 2). In particular, SNPs found in *SCD* (Pauciullo *et al.* 2012a) and

1 *OXT* (Pauciullo *et al.* 2012b) genes were significantly associated with milk production, whereas  
2 recently a polymorphism in the  $\alpha$ s-1 casein encoding gene (*CSN1S1*) was found selectively  
3 neutral in relation to milk yield (Cosenza *et al.* 2014) like observed in the present study for *LTF*  
4 (Insert table 2 here). However, this result does not exclude the existence of a association of *LTF*  
5 polymorphisms in other buffalo populations. In fact, different breeding conditions might have an  
6 indirect effect on milk production, as well as linkage disequilibrium conditions of *LTF* with other  
7 genes influencing milk production traits might result in significant associations with milk yield.  
8 In this context, the lactoferrin gene in cattle is mapped on chromosome 22q24 at a distance of  
9 6.5cM from a QTL for SCS (Heyen *et al.* 1999), whose content was demonstrated to influence  
10 the milk yield and its technological properties (Schutz 1994; Ikonen *et al.* 2004; Cassandro *et al.*  
11 2008). Furthermore, in the same chromosomal region Ashwell *et al.* (2004) identified a QTL  
12 affecting milk production, protein percentage and SCS. In support of this, the buffalo lactoferrin  
13 was already positively correlated with total proteins and SCS and negatively correlated with  
14 lactose and titratable acidity (Giacinti *et al.* 2013), analogously to what was observed for cattle  
15 (Kaminski *et al.* 2006; Parland *et al.* 2010). Therefore further studies are necessary to better  
16 verify this link also in a Mediterranean buffalo population as well in other buffalo breeds.

17 In cattle, *LTF* polymorphisms were also associated with other traits, including  
18 reproductive performances (O'Halloran *et al.* 2010; Zabolewicz *et al.* 2012). Although it is  
19 known that in dairy cows the intense selection for milk production resulted in health and fertility  
20 problems, the genetic progress for this trait is the major challenge also in river buffalo. Therefore,  
21 the improvement of the reproductive performances could be in the next future also a goal for  
22 buffalo selective breeding programmes. In this perspective, the detected polymorphisms at *LTF*  
23 locus can represent a useful investigative tool for future association studies. In fact, the protective  
24 effects of the correlation with SCS or total milk proteins (Giacinti *et al.* 2013) and the potential

1 role in fertility preservation (O'Halloran *et al.* 2010) could be combined to preserve both  
2 productive performances and health of the animals.

#### 4 **Conclusions**

5 The polymorphisms detected in the present work represent the first genetic markers for the  
6 Mediterranean river buffalo lactoferrin encoding gene since no information were available so far  
7 for this breed. The DNA fragment between the exon 15 and 16 resulted extremely polymorphic  
8 considering that a total of 11 SNP were found in about 1500 bp.

9 No association between the exonic SNPs g.88G>A and g.1351G>A and milk yield was  
10 revealed in the present study. Although this result suggests that these polymorphisms are  
11 selectively neutral in relation to the improvement of milk production in Mediterranean river  
12 buffalo reared in Campania, they add important information in terms of genetic variability and  
13 they could be used for future studies of association with other traits of economic interest like  
14 protein and fat production, SCS or reproductive traits.

15 Furthermore, considering that the genome wide SNPchip is currently not available for  
16 river buffaloes, these SNPs might be considered as potential candidate for the development of  
17 this tool and therefore contribute in future genome wide association studies (GWAS).

#### 19 **Acknowledgements**

20 This research was financially supported by the Italian Ministry for Agriculture and  
21 Forestry Policy-MiPAAF (INNOVAGEN project) and by POR CAMPANIA FSE  
22 2007/2013. Sviluppo di Reti di Eccellenza tra Università – Centri di Ricerca – Imprese.  
23 PROGETTO CARINA “Sicurezza, sostenibilità e competitività nelle produzioni  
24 agroalimentari della Campania” CUP B25B09000080007

## Conflict of interest

The authors declare that they have no conflict of interest.

## References

- Arnould VM, Soyeurt H, Gengler N, Colinet FG, Georges MV, Bertozzi C, Portetelle D, Renaville R (2009) Genetic analysis of lactoferrin content in bovine milk. *Journal of Dairy Science* **92**, 2151-2158. doi:10.3168/jds.2008-1255
- Ashwell MS, Heyen DW, Sonstegard TS, Van Tassel CP, Da Y, Van Raden PM, Ron M, Weller JI, Lewin HA (2004) Detection of quantitative trait loci affecting milk production, health, and reproduction traits in Holstein cattle. *Journal of Dairy Science* **87**, 468-475. [http://dx.doi.org/10.3168/jds.S0022-0302\(04\)73186-0](http://dx.doi.org/10.3168/jds.S0022-0302(04)73186-0)
- Bonfatti V, Giantin M, Gervaso M, Rostellato R, Coletta A, Dacasto M, Carnier P (2012) *CSN1S1-CSN3* ( $\alpha$ s1-k-casein) composite genotypes affect detailed milk protein composition of Mediterranean water buffalo. *Journal of Dairy Science* **95**, 6801–6805. <http://dx.doi.org/10.3168/jds.2012-5601>
- Borghese A (2005) Buffalo production and research. Edit by Antonio Borghese, FAO Regional Office for Europe, REU Technical Series, 67
- Cassandro M, Comin A, Ojala M, Dal Zotto R, De Marchi R, Gallo L, Carnier P, Bittante G (2008) Genetic parameters of milk coagulation properties and their relationships with milk yield and quality traits in Italian Holstein cows. *Journal of Dairy Science* **91**, 371-376. <http://dx.doi.org/10.3168/jds.2007-0308>

- 1 Cosenza G, Pauciullo A, Mancusi A, Nicodemo D, Di Palo R, Zicarelli L, Di Berardino D,  
2 Ramunno L (2007) Mediterranean river buffalo oxytocin-neurophysin I (OXT) gene:  
3 structure, promoter analysis and allele detection. *Italian Journal of Animal Science* **6**,  
4 303-306. <http://dx.doi.org/10.4081/ijas.2007.s2.303>
- 5 Cosenza G, Feligini M, Mancusi A, D'Avino A, Coletta A, Di Berardino D, Ramunno L (2009a)  
6 Italian Mediterranean river buffalo CSN2 gene structure and promoter analysis Italian.  
7 *Journal of Animal Science* **8**, 57-59. <http://dx.doi.org/10.4081/ijas.2009.s2.57>
- 8 Cosenza G, Pauciullo A, Feligini M, Coletta A, Colimoro L, Di Berardino D, Ramunno L  
9 (2009b) A point mutation in the splice donor site of intron 7 in the  $\alpha$ s2-casein encoding  
10 gene of the Mediterranean River buffalo results in an allele-specific exon skipping.  
11 *Animal Genetics* **40**, 791. doi: 10.1111/j.1365-2052.2009.01897.x
- 12 Cosenza G, Pauciullo A, Macciotta NPP, Apicella E, Steri R, La Battaglia A, Jemma L, Coletta  
13 A, Di Berardino D, Ramunno L (2014) Mediterranean River Buffalo *CSN1S1* gene:  
14 search for polymorphisms and association studies. *Animal Production Science*, accepted  
15 16 February. <http://dx.doi.org/10.1071/AN13438>
- 16 Daly M, Ross P, Giblin L, Buckley F (2006) Polymorphisms within the lactoferrin gene  
17 promoter in various cattle breeds *Animal Biotechnology* **17**, 33–42. doi:  
18 10.1080/10495390500460965
- 19 Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, Altshuler D  
20 (2002) The structure of haplotype blocks in the human genome. *Science* **296**, 2225–2229.  
21 doi: 10.1126/science.1069424
- 22 Giacinti G, Basiricò L, Ronchi B, Bernabucci U (2013) Lactoferrin concentration in buffalo milk.  
23 *Italian Journal of Animal Science* **12:e23**, 139-143.  
24 <http://dx.doi.org/10.4081/ijas.2013.e23>



- 1 Goossens M, Kan YW (1981) DNA analysis in the diagnosis of hemoglobin disorders. *Methods*  
2 *in Enzymology* **76**, 805-817. [http://dx.doi.org/10.1016/0076-6879\(81\)76159-7](http://dx.doi.org/10.1016/0076-6879(81)76159-7)
- 3 Guo BL, Jiao Y, He C, Wei LX, Chang ZH, Yue XP, Lan XY, Chen H, Lei CZ (2010) A novel  
4 polymorphism of the lactoferrin gene and its association with milk composition and body  
5 traits in dairy goats. *Genetics and Molecular Research* **9**, 2199-2206. doi: 10.4238/vol9-  
6 4gmr928
- 7 Heinemeyer T, Wingender E, Reuter I, Hermjakob H, Kel AE, Kel OV, Ignatieva EV, Ananko  
8 EA, Podkolodnaya OA, Kolpakov FA, Podkolodny NL, Kolchanov NA (1998) Databases  
9 on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acids*  
10 *Research* **26**, 362-367. doi: 10.1093/nar/26.1.362
- 11 Heyen DW, Weller JI, Ron M, Band M, Beever JE, Feldmesser E, Da Y, Wiggans GR,  
12 VanRaden PM, Lewin HA (1999) A genome scan for QTL influencing milk production  
13 and health traits in dairy cattle. *Physiological Genomics* **1**, 165-175.  
14 <http://physiolgenomics.physiology.org/content/1/3/165>
- 15 Huang J, Wang H, Wang C, Li J, Li Q, Hou M, Zhong J (2010). Single nucleotide  
16 polymorphisms, haplotypes and combined genotypes of lactoferrin gene and their  
17 associations with mastitis in Chinese Holstein cattle. *Molecular Biology Reports* **37**, 477–  
18 483. doi: 10.1007/s11033-009-9669-1
- 19 Ikonen T, Morri A, Tyriseva AM, Ruottinen O, Ojala M (2004) Genetic and phenotypic  
20 correlations between milk coagulation properties, milk production traits, somatic cell  
21 count, casein content and pH of milk. *Journal of Dairy Science* **87**, 458-467.  
22 [http://dx.doi.org/10.3168/jds.S0022-0302\(04\)73185-9](http://dx.doi.org/10.3168/jds.S0022-0302(04)73185-9)

- 1 Kang JF, Li XL, Zhou RY, Li LH, Feng FJ, Guo XL (2008) Bioinformatics analysis of  
2 lactoferrin gene for several species. *Biochemical Genetics* **46**, 312-322. doi:  
3 10.1007/s10528-008-9147-9
- 4 Kaminski S, Olenski K, Brym P, Malewski T, Sazanov AA (2006) Single Nucleotide  
5 Polymorphism in the promoter region of the lactoferrin gene and its associations with  
6 milk performance traits in Polish holstein–friesian cows. *Russian Journal of Genetics* **4**,  
7 924-927. doi: 10.1134/S1022795406080114
- 8 Kathiravan P, Kataria RS, Mishra BP (2009) Single nucleotide polymorphism (SNP)  
9 identification and sequence analysis of 5' flanking region of lactoferrin gene in Indian  
10 buffaloes (Bubalus bubalis). *Livestock Science* **121**, 38-44.  
11 <http://dx.doi.org/10.1016/j.livsci.2008.05.021>
- 12 Kathiravan P, Kataria RS, Mishra BP, Dubey PK, Selvakumar M, Tyagi N (2010) Seven novel  
13 single nucleotide polymorphisms identified within river buffalo (Bubalus bubalis)  
14 lactoferrin gene. *Tropical Animal Health and Production* **42**, 1021-1026. doi:  
15 10.1007/s11250-009-9516-3
- 16 Khatibi M, Roshanfekr H, Fayazi J, Mirzade K (2013) Polymorphism of 5' flanking region of  
17 lactoferrin gene in Khuzestan buffaloes. *International Journal of Advanced Biological*  
18 *and Biomedical Research* **1**, 777-782.
- 19 Li GH, Zhang Y, Sun DX, Li N (2004) Study on the polymorphism of bovine lactoferrin gene  
20 and its relationship with mastitis. *Animal Biotechnology* **15**, 67–76. doi: 10.1081/ABIO-  
21 120037899
- 22 Lee TH, Shimazaki K, Yu SL, Nam MS, Kim SJ, Lee KK, Yu DY (1997) Polymorphic sequence  
23 of Korean native goat lactoferrin exhibiting greater antibacterial activity. *Animal Genetics*  
24 **28**, 367–369. doi: 10.1111/j.1365-2052.1997.00154.x

- 1 Masina P, Rando A, Di Gregorio P, Cosenza G, Mancusi A (2007) Water buffalo kappa-casein  
2 gene sequence. *Italian Journal of Animal Science* **6**, 353-355. doi:  
3 10.4081/ijas.2007.s2.353
- 4 O'Halloran F, Bahar B, Buckley F, O'Sullivan O, Sweeney T, Giblin L (2009) Characterisation  
5 of single nucleotide polymorphisms identified in the bovine lactoferrin gene sequences  
6 across a range of dairy cow breeds. *Biochimie* **91**, 68-75. doi:  
7 10.1016/j.biochi.2008.05.011
- 8 O'Halloran F, Berry DP, Bahar B, Howard DJ, Sweeney T, Giblin L (2010) Polymorphisms in  
9 the bovine lactoferrin promoter are associated with reproductive performance and somatic  
10 cell count. *Journal of Dairy Science* **93**, 1253-1259. doi: 10.3168/jds.2009-2699
- 11 Parland Mc, Giblin L, Veerkamp RF, Berry DP (2010) The impact of selection on milk  
12 production on the lactoferrin content of milk in Irish Holstein-Friesians. *Advances in*  
13 *Animal Biosciences*, Proceedings of the British Society of Animal Science and the  
14 Agricultural Research Forum **1**, 305. <http://dx.doi.org/10.1017/S2040470010004486>
- 15 Pauciullo A, Cosenza G, D'Avino A, Colimoro L, Nicodemo D, Coletta A, Feligini M,  
16 Marchitelli C, Di Berardino D, Ramunno L (2010a) Sequence analysis and genetic  
17 variability of stearoyl CoA desaturase (SCD) gene in the Italian Mediterranean river  
18 buffalo. *Molecular and Cellular Probes* **24**, 407-410. doi: 10.1016/j.mcp.2010.07.009
- 19 Pauciullo A, Cosenza G, Nicodemo D, Gallo D, Mancusi A, Crepaldi P, Di Berardino D,  
20 Ramunno L (2010b) Molecular cloning, promoter analysis and SNP identification of  
21 Italian Nicastrrese and Saanen lactoferrin gene. *Veterinary Immunology and*  
22 *Immunopathology* **134**, 279-283. doi: 10.1016/j.vetimm.2009.09.002

- 1 Pauciullo A, Cosenza G, Di Berardino D, Ramunno L (2011) Single Nucleotide Polymorphism at  
2 the LGB gene of the Italian Mediterranean River Buffalo. *Deutschen Gesellschaft für*  
3 *Züchtungskunde* (DGfZ), 6-7 September, Freising - Weihenstephaan, pp. A06
- 4 Pauciullo A, Cosenza G, Steri R, Coletta A, La Battaglia A, Di Berardino D, Macciotta NPP,  
5 Ramunno L (2012a) A single nucleotide polymorphism (SNP) in the promoter region of  
6 River buffalo Stearoyl CoA Desaturase gene (SCD) is associated with milk yield. *Journal*  
7 *of Dairy Research* **79**, 429-435. <http://dx.doi.org/10.1017/S0022029912000507>
- 8 Pauciullo A, Cosenza G, Steri R, Coletta A, Jemma L, Feligini M, Di Berardino D, Macciotta  
9 NPP, Ramunno L (2012b) An association between Single Nucleotide Polymorphisms at  
10 the Oxytocin locus and milk yield in Italian Mediterranean river buffalo. *Journal of Dairy*  
11 *Research* **79**, 150-156. doi: 10.1017/S0022029911000914
- 12 Risch NJ (2000) Searching for genetic determinants in the new millennium. *Nature* **405**, 847-856.  
13 doi:10.1038/35015718
- 14 Schutz MM (1994) Genetic evaluation of Somatic Cell Scores for United States dairy cattle  
15 *Journal of Dairy Science* **77**, 2113-2129. doi:10.3168/jds.S0022-0302(94)77154-X
- 16 Seyfert HM, Tuckoricz A, Interthal H, Koczan D, Hobom G (1994) Structure of the bovine  
17 lactoferrin encoding gene and its promoter. *Gene* **143**, 265-269.  
18 [http://dx.doi.org/10.1016/0378-1119\(94\)90108-2](http://dx.doi.org/10.1016/0378-1119(94)90108-2)
- 19 Vogel HJ, Schibli D, Jing W, Lohmeier-Vogel EM, Epand RF, Epand RM (2002) Towards a  
20 structure function analysis of bovine lactoferricin and related tryptophan and arginine  
21 containing peptides. *Biochemistry and Cell Biology* **80**, 49-63. doi: 10.1139/o11-057
- 22 Zabolewicz T, Brym P, Olenski K, Suchocki K, Malewski T, Szyda J, Kaminski S (2012)  
23 Polymorphism within TATA-box of bovine lactoferrin gene and its association with

1 performance traits in Holstein cattle. *Livestock Science* **149**, 267–274.  
2 <http://dx.doi.org/10.1016/j.livsci.2012.07.020>  
3 Wang M, Moisés S, Khan MJ, Wang J, Bu D, Loo JJ (2012) MicroRNA expression patterns in  
4 the bovine mammary gland are affected by stage of lactation. *Journal of Dairy Science*  
5 **95**, 6529–6535. doi: 10.3168/jds.2012-5748

Table 1. Genotyping data, allele frequency, absolute (relative) frequencies of buffalo cows, lactations and tests across genotypes of the SNPs:  
g.88G>A SNP at the exon 15 and g.1351G>A at the exon 16 of the *LTF* gene in Mediterranean river buffalo population

Position		Genotype distribution*				Allele freq.*		Absolute and relative frequencies used in the model			
								Genotype	n of cows	n of lactations	N of tests
Exon 15/Exon16		aa	ab	bb	TOT	a	b	aa	253 (45,75)	501 (45,67)	3,268 (45,01)
	Obs.	253	248	52	553	0,68	0,32	ab	248 (44,84)	488 (44,48)	3,278 (45,15)
	Exp.	257,01	239,97	56,01				bb	52 (9,41)	108 (9,85)	714 (9,84)
	$\chi^2=0,619$							total	553 (100)	1097 (100)	7,260 (100)

\* allele a = G and allele b = A for the SNP g.88G>A; allele a = A and allele b = G for the SNP g.1351G>A

Table 2. Comparison among the least squares means of milk yield (kg/d) for the three genotypes at the *LTF* g.88G>A SNP estimated with the model (1) and the milk yield data (kg/d) reported for *SCD* (Pauciullo *et al.* 2012a), *OXT* (Pauciullo *et al.* 2012b), and *CSN1S1* (Cosenza *et al.* 2014).

<i>LTF</i>	MY	SE	<i>SCD</i>	MY	<i>OXT</i>	MY	<i>CSN1S1</i>	MY
GG	8.35	0.08	AA	8.63 <sup>ab</sup>	CC	8.73 <sup>ab</sup>	CC	7.81
AG	8.19	0.08	CA	8.83 <sup>a</sup>	CT	8.21 <sup>a</sup>	CT	7.92
AA	8.55	0.15	CC	6.60 <sup>b</sup>	TT	10.07 <sup>b</sup>	TT	7.56

<sup>a,b</sup> Means within columns with different superscripts differ (Bonferroni adjusted  $P < 0.05$ )




Fig 1. Haplotypes of the *LTF* gene in river buffalo population. Standard color scheme is used to display linkage disequilibrium. Numbers next to each haplotype bar are haplotype frequencies. On top of the figure, a schematic representation of the sequenced regions of the river buffalo LTF gene. Exons are indicated by boxes whereas the intron 15 is showed by an unbroken line.